

INFINITE DIMENSIONAL GEOMETRY AND
QUANTUM FIELD THEORY OF STRINGS. III
INFINITE DIMENSIONAL W-GEOMETRY OF
A SECOND QUANTIZED FREE STRING.

D.V. JURIEV¹

*Laboratoire de Physique Théorique
de l'École Normale Supérieure,
24 rue Lhomond, 75231 Paris Cedex 05, France²
E mail: juriev@physique.ens.fr*

ABSTRACT. The present paper is devoted to various objects of the infinite dimensional W -geometry of a second quantized free string. Our purpose is to include the W -symmetries into the general infinite dimensional geometrical picture related to the quantum field theory of strings, which was described in [1]. It is done by a change of the Lie algebra of all infinitesimal reparametrizations of a string world-sheet on the Lie quasi(pseudo)algebra of classical W -transformations (Gervais-Matsuo quasi(pseudo)algebra) as well as of the Virasoro algebra on the central extended enlarged Gervais-Matsuo quasi(pseudo)algebra. A way to obtain W -algebras from the classical W -transformations (i.e. Gervais-Matsuo Lie quasi(pseudo)algebra) is proposed. The relations of Gervais-Matsuo differential W -geometry to the Batalin-Weinstein-Karasev-Maslov approach to the geometry of nonlinear Poisson brackets as well as to L.V.Sabinin program of "nonlinear geometric algebra" are mentioned.

Keywords: Infinite dimensional geometry, string field theory, W-symmetries and W-geometry, Lie quasi(pseudo)algebras.

1991 MSC: 58B25, 17B65, 81T30, 17D99

INTRODUCTION

This is the third part of the text devoted to an infinite dimensional geometry of a quantum field theory of strings (for previous ones see [1,2]). It is devoted to an infinite dimensional geometry, which appears in the theory of second quantized free strings, as well as the first one. An origin of the writing of this article is hidden in some very crucial detail of the theory, which is known under the title of W -symmetries. The algebraic aspects of W -symmetries, cumbersome

¹On leave from Mathematical Division of Research Institute for System Studies, Moscow, Russia (e mail: juriev@systud.msk.su)

²Unité Propre du Centre National de la Recherche Scientifique associée à l'École Normale Supérieure et à l'Université de Paris-Sud

and mysterious a little at the beginning [3], were recently clarified by B.A.Khesin, V.Yu.Ovsienko, A.O.Radul et al [4-10]. Moreover, a simple geometric interpretation of them was given by J.-L.Gervais and Y.Matsuo [11]. It seems that the cruciality of W -symmetries to the theory of second quantized free strings is in the fact that W -”reparametrizations” of strings, which are considered, may be not obligatory global in the loop space (i.e. unique for all strings independently on their position in a target space) but in some sense local, i.e. depends in some natural way on an embedding of the string into the target space. One may say that such transformations are related not only to the intrinsic but also extrinsic geometry of a string. A class of the most interesting transformations of this kind, which are related to a certain complex analog of a classical Frenet theory of curves invariants [12], was described by J.-L.Gervais and Y.Matsuo as classical W -transformations. Such transformations of both internal and external degrees of freedom of a string maybe considered as hidden ones with respect to standard conformal symmetries. So a natural question arises: how the process of a geometric quantization of a string changes under an account of these hidden symmetries. From the mathematical point of view it may be considered as a problem of an ”induction” of the quantization process directed by a considered symmetry algebra to one directed by its extension. So the solution of this problem (reproduced from a typical pattern for this partial infinite dimensional case of the quantum field theory of free strings, which we are considered) should produce a lot of new geometric material, completing a general picture, which was drawn in [1], by new intriguing details and nuances.

This opinion is also confirmed by the fact that W_∞ -symmetries are very deeply related to the group of area-preserving diffeomorphisms of a torus. This group, the corresponding Lie algebra, its deformations and central extensions, being the main objects of a certain development of the string theory, the membrane theory, produce the next scope of infinite dimensional objects after one of the Virasoro algebra. So we may consider a geometric material, which will be discussed in the paper, as a possible introduction to the transition from the careful consideration of the infinite dimensional geometry of objects from the ”Virasoro family”, which was begun by A.A.Kirillov and the author in [13] and summarized partially in [14,1,2] (see also references wherein) to a systematic treatment of geometric material, concerning objects related to the group of symplectomorphisms of a torus.

Such treatment is supposed to be presented in one of the forthcoming publications.

In the aspect of the applications of our geometric material we follow the general ideology of A.Yu.Morozov, formulated in [15], which proposed to consider the string theory not as a partial physical theory describing some narrow class of real phenomena but as one of the universal theories, which final and, it seems, rather wide area of applications is not understood completely now. It is possible that some new applications of certain aspects of the string theory (and of an infinite dimensional geometry related to it) will be found during the performing of an interdisciplinary ”toy-program” of an investigation of peculiarities of human vision (in particular, of color perception) in artificial interactive computerographical systems, formulated by the author in [16], the results of which may be important for understanding of various processes in natural interactive (sensorial and visual) systems. Hence, our presentation is maximally mathematical and avoids many technical aspects, which are crucial for some applications; instead of that we prefer to specially point out a general mathematical meaning of our constructions.

The author thanks Prof.J.-L.Gervais and the collective of Laboratoire de Physique Théorique de l'École Normale Supérieure (Paris) for a friendly imaginative, creative atmosphere and hospitality, without which the writing of this paper will be impossible. The author also specially thanks Prof.J.-L.Gervais, whose ideas on differential W-geometry provoked him to realize a plan of the article. Also the author is undebtful to all participants of the seminar on quantum field theory (especially, Prof.A.Yu.Morozov and A.O.Radul) at the Institute for Theoretical and Experimental Physics (Moscow), on which the problems related to W-symmetries were discussed for many times. The author thanks M.I.Golenishcheva-Kutuzova, B.A.Khesin and V.Yu.Ovsienko, who being participants of the seminar on representation theory at Department of Mathematics of Moscow State University introduced the author into the theme, Prof.M.V.Saveliev (Institute of High Energy Physics, Serpukhov) for some conversations on differential W-geometry at LPTENS, Prof.L.V.Sabinin and P.O.Mikheev (Dpt.Mathematics, Univ.Peoples' Friendship, Moscow) for discussions on Lie quasi(pseudo)algebras, Lie quasi(pseudo)groups and related topics.

1. SYMPLECTOMORPHISMS (AREA-PRESERVING TRANSFORMATIONS) OF TORUS, DIFFERENTIAL OPERATORS ON A CIRCLE AND W -ALGEBRAS

1.1. Groups $\text{Symp}(\mathbb{T}^2)$, $\text{Symp}^e(\mathbb{T}^2)$ and $\text{Symp}_0(\mathbb{T}^2)$ of symplectomorphisms of a torus $\mathbb{T}^2$ and their Lie algebras $\text{Ham}(\mathbb{T}^2)$ and $\text{Ham}_0(\mathbb{T}^2)$ of Hamiltonian and strictly Hamiltonian vector field on a torus. Poisson algebras $\mathcal{F}^{\mathbb{C}}(\mathbb{T}^2)$, $\mathcal{F}^{\mathbb{C}}(T^*(S^1))$ and $\mathcal{F}^{\mathbb{C}}(\mathbb{R}^2)$ of functions on tori, cylinder and plane, their deformations and central extensions.

Let $(\mathbb{T}^2, \omega)$ be a two-dimensional torus with a fixed 2-form (a volume form) ω on it. Let $\text{Symp}(\mathbb{T}^2, \omega)$ be a group of symplectomorphisms of a torus $(\mathbb{T}^2, \omega)$, i.e. all diffeomorphisms ζ of $\mathbb{T}^2$ such that $\zeta_*\omega = \omega$ [17]. So symplectomorphisms of a torus are just *its area-preserving transformations*.

It may be easily shown that all groups $\text{Symp}(\mathbb{T}^2, \omega)$ are isomorphic. Indeed, let's identify a torus $(\mathbb{T}^2, \omega)$ with the quotient $\mathbb{R}^2/\Gamma$, where the plane $\mathbb{R}^2$ with a fixed coordinate system (p, q) possesses a canonical 2-form $dp \wedge dq$, Γ is a certain free lattice on $\mathbb{R}^2$, i.e. a set $\mathbb{Z}\mathbf{a} + \mathbb{Z}\mathbf{b}$, where $\mathbf{a}$ and $\mathbf{b}$ are two independent vectors on the plane. The symplectomorphisms of a torus may be identified with the classes mod Γ of functions $\mathbf{f} : \mathbb{R}^2 \mapsto \mathbb{R}^2$, which obey some conditions. The expression mod Γ means that two functions $\mathbf{f}_1$ and $\mathbf{f}_2$ are called equivalent iff $\mathbf{f}_1(\mathbf{x}) - \mathbf{f}_2(\mathbf{x}) \in \Gamma$ for all $\mathbf{x}$ from $\mathbb{R}^2$. It should be mentioned that in the coordinate form a function $\mathbf{f}$ is represented as a pair (f_p, f_q) of functions $f_p = f_p(p, q)$ and $f_q = f_q(p, q)$. The conditions, to which a function $\mathbf{f}$ should obey, are the next ones: (1) the periodicity condition: $\mathbf{f}(\mathbf{x} + m\mathbf{a} + n\mathbf{b}) = \mathbf{f}(\mathbf{x}) + m\tilde{\mathbf{a}} + n\tilde{\mathbf{b}}$ for all $\mathbf{x}$ from $\mathbb{R}^2$, where $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$ is a pair of generators of Γ , i.e. $\tilde{\mathbf{a}} = A(\mathbf{a})$, $\tilde{\mathbf{b}} = A(\mathbf{b})$, A is a certain transformation from $\text{Aut}(\Gamma)$, which is isomorphic to $\text{SL}(2, \mathbb{Z})$; (2) the normalization of Jacobian condition: $\frac{D\mathbf{f}}{D\mathbf{x}} = \det \frac{\partial(f_p, f_q)}{\partial(p, q)} = 1$. The last determinant (Jacobian) is invariant under all linear transformations of a plane $\mathbb{R}^2$. Because for arbitrary two free lattices Γ_1 and Γ_2 there exists a linear transformation γ of $\mathbb{R}^2$ such that $\Gamma_2 = \gamma(\Gamma_1)$, the group of symplectomorphisms of $\mathbb{R}^2/\Gamma_1$ and $\mathbb{R}^2/\Gamma_2$ may be identified. So instead of the notation $\text{Symp}(\mathbb{T}^2, \omega)$ for a group of symplectomorphisms of a torus we shall use the simpler notation $\text{Symp}(\mathbb{T}^2)$.

It should be mentioned that the group $\text{Symp}(\mathbb{T}^2)$ is not connected. Let us denote the component of the identity in it by $\text{Symp}^e(\mathbb{T}^2)$. Of course, $\text{Symp}^e(\mathbb{T}^2)$ is a normal subgroup in $\text{Symp}(\mathbb{T}^2)$ and therefore we have the following strict sequence:

$$0 \longrightarrow \text{Symp}^e(\mathbb{T}^2) \longrightarrow \text{Symp}(\mathbb{T}^2) \longrightarrow \text{Symp}(\mathbb{T}^2)/\text{Symp}^e(\mathbb{T}^2) \longrightarrow 0.$$

The quotient $\text{Symp}(\mathbb{T}^2)/\text{Symp}^e(\mathbb{T}^2)$ is isomorphic to *the modular group* $\text{SL}(2, \mathbb{Z})$, so that the functions $\mathbf{f}$ corresponded to the elements of $\text{Symp}^e(\mathbb{T}^2)$ obey the periodicity condition in the form $\mathbf{f}(\mathbf{x} + m\mathbf{a} + n\mathbf{b}) = \mathbf{f}(\mathbf{x}) + m\mathbf{a} + n\mathbf{b}$.

The group $\text{Symp}^e(\mathbb{T}^2)$ admits a normal subgroup $\text{Symp}_0(\mathbb{T}^2)$ of such symplectomorphisms ζ that $\int_{-1/2}^{1/2} \int_{-1/2}^{1/2} \mathbf{f}(\mathbf{x} + t_1\mathbf{a} + t_2\mathbf{b}) dt_1 dt_2 = \mathbf{x}$. The quotient $\text{Symp}^e(\mathbb{T}^2)/\text{Symp}_0(\mathbb{T}^2)$ is isomorphic to 2-dimensional compact abelian group $\mathbb{T}^2$, so that a strict sequence

$$0 \longrightarrow \text{Symp}_0(\mathbb{T}^2) \longrightarrow \text{Symp}^e(\mathbb{T}^2) \longrightarrow \mathbb{T}^2 \longrightarrow 0$$

exists. This strict sequence may be resolved, the resolving mapping $\mathbb{T}^2 \rightarrow \text{Symp}^e(\mathbb{T}^2)$ realises a group $\mathbb{T}^2$ as a group of movements of a torus $\mathbb{T}^2$.

The Lie algebra of the group $\text{Symp}^e(\mathbb{T}^2)$ is *an algebra of Hamiltonian vector fields on a torus*, i.e. the fields ξ such that $\mathcal{L}_\xi \omega = 0$. It should be mentioned that after an identification of a torus $\mathbb{T}^2$ with the quotient $\mathbb{R}^2/\Gamma$ Hamiltonian fields ξ on a torus may be also characterized as divergence free ones, i.e. vector fields ξ such that $\text{div } \xi = 0$. From Cartan theorem it follows that ξ is Hamiltonian iff $d\alpha_\xi = 0$, where $\alpha_\xi = \iota_\xi \omega$. So the algebra $\text{Ham}(\mathbb{T}^2)$ of Hamiltonian vector fields on a torus may be identified with an algebra of closed 1-forms on a torus with respect to the bracket $\{\alpha, \beta\} = d\frac{\alpha \wedge \beta}{\omega}$.

The Lie algebra of the group $\text{Symp}_0(\mathbb{T}^2)$ is a certain subalgebra of $\text{Symp}^e(\mathbb{T}^2)$, *the algebra of strictly Hamiltonian vector fields*, i.e. the Hamiltonian vector fields ξ such that the corresponding 1-form α_ξ is strict. The Lie algebra $\text{Ham}_0(\mathbb{T}^2)$ of strictly Hamiltonian vector fields on a torus is an ideal in the algebra $\text{Ham}(\mathbb{T}^2)$ so that a strict sequence

$$0 \longrightarrow \text{Ham}_0(\mathbb{T}^2) \longrightarrow \text{Ham}(\mathbb{T}^2) \longrightarrow \mathbb{R}^2 \longrightarrow 0$$

exists. The quotient $\mathbb{R}^2 = \text{Ham}(\mathbb{T}^2)/\text{Ham}_0(\mathbb{T}^2)$ may be interpreted as the cohomology group $H^1(\mathbb{T}^2, \mathbb{R})$ if one considers its as a quotient of a space of closed 1-forms by a subspace of strict ones. This strict sequence may be resolved. The resolving mapping $\mathbb{R}^2 \mapsto \text{Ham}(\mathbb{T}^2)$ realises the vector space $\mathbb{R}^2$ as a certain parallelization of a torus.

It should be mentioned that if 1-form α_ξ is strict then there exist a function F_ξ on a torus such that $\alpha_\xi = dF_\xi$. The bracket in the space of strictly Hamiltonian vector fields induces a bracket in the space $\mathcal{F}(\mathbb{T}^2)$ of functions (0-forms) on a torus. The bracket in the space $\mathcal{F}(\mathbb{T}^2)$ has the form $\{F, G\} = \frac{dF \wedge dG}{\omega}$. The center of the Lie algebra $\mathcal{F}(\mathbb{T}^2)$ consists of constant functions, so it may be identified with 1-dimensional space $\mathbb{R}$. The Lie algebra of strictly Hamiltonian vector fields on a torus is isomorphic to a quotient $\mathcal{F}(\mathbb{T}^2)/\mathbb{R}$. The isomorphism is realised by an exterior derivative d , which maps the above quotient onto the Lie algebra of strict 1-forms, which is isomorphic to $\text{Ham}_0(\mathbb{T}^2)$. The mapping from the algebra $\mathcal{F}(\mathbb{T}^2)$ to the algebra $\text{Ham}_0(\mathbb{T}^2)$ is symbolized as sgrad and is called by "*skew-gradient*",

so that $\xi = \text{sgrad}(F_\xi)$ for all strictly Hamiltonian vector fields on a torus. It should be mentioned that a strict sequence

$$0 \longrightarrow \mathbb{R} \longrightarrow \mathcal{F}(\mathbb{T}^2) \longrightarrow \text{Ham}_0(\mathbb{T}^2) \longrightarrow 0$$

may be resolved. The resolving mapping identify the algebra $\text{Ham}_0(\mathbb{T}^2)$ with the subspace $\mathcal{F}_0(\mathbb{T}^2)$ of functions of zero mean value.

It should be mention that the bracket $\{\cdot, \cdot\}$ on the space $\mathcal{F}(\mathbb{T}^2)$ of functions on a torus, determines a structure of a Poisson algebra on it. This algebra is called by *the Poisson algebra of functions on a torus*. It means that $\{F, GH\} = \{F, G\}H + G\{F, H\}$ for all functions F, G, H from $\mathcal{F}(\mathbb{T}^2)$. In another words the bracket $\{\cdot, \cdot\}$ is a derivation of the ordinary commutative multiplication of functions.

Let us realise torus $\mathbb{T}^2$ as a quotient $\mathbb{R}^2/\Gamma$, where $\Gamma = \{(mM, nN) \in \mathbb{R}^2, m, n \in \mathbb{Z}; M, N \text{ are fixed real positive numbers}\}$. Then there is defined a canonical basis in $\mathcal{F}^\mathbb{C}(\mathbb{T}^2)$, the complexification of the algebra $\mathcal{F}(\mathbb{T}^2)$: $e_{m,n} = \mathcal{D}(M, N) \exp(\frac{im\pi}{M}) \exp(\frac{inq}{N})$, where it is convenient to put a normalization constant $\mathcal{D}(M, N)$ being equal to MN . The Poisson brackets $\{\cdot, \cdot\}$ in $\mathcal{F}(\mathbb{T}^2)$ or $\mathcal{F}^\mathbb{C}(\mathbb{T}^2)$ have the form $\{F, G\} = \frac{\partial F}{\partial p} \frac{\partial G}{\partial q} - \frac{\partial G}{\partial p} \frac{\partial F}{\partial q}$ so that [17]

$$\{e_{m,n}, e_{m',n'}\} = (mn' - nm')e_{m+m', n+n'}.$$

The subalgebra $\mathcal{F}_0^\mathbb{C}(\mathbb{T}^2)$ of $\mathcal{F}^\mathbb{C}(\mathbb{T}^2)$ (the complexification of $\mathcal{F}_0(\mathbb{T}^2)$) is realized as one spanned by generators $e_{m,n}, (m, n) \neq (0, 0)$.

Let us call the algebra spanned formally by $e_{m,n}$ by *Floratos-Iliopoulos algebra* (see [18], where it was considered). It should be mentioned that generators $L_k, k \in \mathbb{Z}$ of the algebra $\mathbb{C}\text{Vect}(\mathbb{S}^1)$ of vector fields on a circle $\mathbb{S}^1$, obeying the commutation relations $[L_i, L_j] = (i - j)L_{i+j}$ may be expressed formally via $e_{m,n}$ in the following way [18]:

$$L_k = \sum_{m \in \mathbb{Z}} \mathcal{G}_k(m) e_{m,k},$$

where $\mathcal{G}_k(m) = \frac{(-1)^m}{m}$ if $m \neq 0$ and 0 otherwise. Of course, under the verification of the commutations relations of generators L_k some rather natural regularization is used. So the Lie algebra $\mathbb{C}\text{Vect}(\mathbb{S}^1)$ (more precisely, so called *Witt algebra*, which is spanned by generators L_k of $\mathbb{C}\text{Vect}(\mathbb{S}^1)$), but we shall not mention a difference between them below) is realized as a certain subalgebra of the Floratos-Iliopoulos algebra.

Under the action of the algebra $\mathbb{C}\text{Vect}(\mathbb{S}^1)$ the Floratos-Iliopoulos algebra may be decomposed into the sum of tensor modules of positive integer weights, the corresponding tensor operators will be denoted by $w_n^{(i)}$ ($i \geq 2$). It should be mentioned that $L_n = w_n^{(2)}$, the commutation relations between generators $w_n^{(i)}$ have the form

$$[w_n^{(i)}, w_m^{(j)}] = ((j-1)n - (i-1)m)w_{n+m}^{(i+j-2)},$$

$w_n^{(i)}$ themselves are defined as

$$w_k^{(i)} = \sum_{m \in \mathbb{Z}} \mathcal{G}_k^{(i)}(m) e_{k,m},$$

$\mathcal{G}_k^{(i)}(m) = P_i(\frac{1}{m})$, where the polynomials $P_i(x)$ of degree $i - 1$ are connected by the easily calculated recurrent formulas.

The algebra $\widehat{\text{Ham}}_0(\mathbb{T}^2)$ of strictly Hamiltonian vector fields on a torus admits a central extension $\widehat{\text{Ham}}_0(\mathbb{T}^2)$, which may be defined in a following way $[\xi, \eta]_{\partial} = [\xi, \eta] + \int_{\mathbb{T}^2} [\partial, \xi] \eta \, dpdq$, where ∂ is a certain Hamiltonian but not strictly Hamiltonian vector field on the torus. One may choose $\partial = a\partial_p + b\partial_q$ and this is a general setting up to trivial central extensions. So the universal central extension of $\text{Ham}_0(\mathbb{T}^2)$ is defined by the strict sequence

$$0 \longrightarrow H_1(\mathbb{T}^2, \mathbb{R}) \longrightarrow \widehat{\text{Ham}}_0(\mathbb{T}^2) \longrightarrow \text{Ham}_0(\mathbb{T}^2) \longrightarrow 0,$$

where $H_1(\mathbb{T}^2, \mathbb{R})$ is a 2-dimensional group of homology of torus with real coefficients.

The central extension $\widehat{\text{Ham}}_0(\mathbb{T}^2)$ of the Lie algebra $\text{Ham}_0(\mathbb{T}^2)$ induces a central extension $\widehat{\mathcal{F}}^{\mathbb{C}}(\mathbb{T}^2)$ of the algebra $\mathcal{F}^{\mathbb{C}}(\mathbb{T}^2)$, defined as $\{F, G\}_{\partial} = \{F, G\} + \int_{\mathbb{T}^2} \mathcal{L}_{\partial} F \cdot G \, dpdq$, so that the following strict sequence

$$0 \longrightarrow \mathbb{C}^2 \longrightarrow \widehat{\mathcal{F}}^{\mathbb{C}}(\mathbb{T}^2) \longrightarrow \mathcal{F}^{\mathbb{C}}(\mathbb{T}^2) \longrightarrow 0$$

exists. This central extension has the following form in the basis $e_{m,n}$:

$$[e_{m,n}, e_{m',n'}] = (mn' - nm')e_{m+m',n+n'} + (am + bn)\delta(m + m')\delta(n + n')\mathbf{1}$$

The extended Floratos-Iliopoulos algebra admits an embedding of the Virasoro algebra $\mathbb{C}\text{vir}$ with a non-trivial central charge into it (one of possible embeddings was described in [18]).

The Poisson algebra $\mathcal{F}^{\mathbb{C}}(\mathbb{T}^2)$ may be deformed into a certain associative algebra [19]. In the canonical basis such deformation has the form

$$e_{m,n} \cdot e_{m',n'} = \exp(i\lambda(mn' - nm'))e_{m+m',n+n'}.$$

The corresponding commutator algebra has the form (after a renormalization of generators)

$$[e_{m,n}, e_{m',n'}] = \frac{1}{\lambda} \sin(\lambda(nm' - mn'))e_{m+m',n+n'}.$$

So there is defined a certain deformation of Floratos-Iliopoulos algebra, which is called (centreless) *Sine-algebra (or Fairlie-Fletcher-Zachos algebra)* [20]. The corresponding deformation of $\text{Ham}_0(\mathbb{T}^2)$ is called *the Lie algebra of strictly Hamiltonian vector fields on a quantum torus* and is denoted by $\text{Ham}_0(\mathbb{T}_q^2)$ ($q = e^{i\lambda}$).

(Centreless) Sine algebra admits a central extension, which has the form

$$[e_{m,n}, e_{m',n'}] = \frac{1}{\lambda} \sin(\lambda(nm' - mn'))e_{m+m',n+n'} + (am + bn)\delta(m + m')\delta(n + n')\mathbf{1}.$$

The Sine-algebra with a center also admits the embedding of the Virasoro algebra $\mathbb{C}\text{vir}$:

$$L_k = \sum_{m \in \mathbb{Z}} \tilde{\mathcal{G}}_k(m) e_{k,m},$$

where $\tilde{\mathcal{G}}_k(m)$ are defined by formulas, analogous to one of [18].

Under the action of the algebra $\mathbb{C}\text{vir}$ Sine-algebra may be also decomposed into the sum of tensor modules of positive integer weights, the corresponding tensor operators will be denoted by $\tilde{w}_n^{(i)}$ ($i \geq 2$). Their commutation relations have the form (for the centreless Sine-algebra)

$$[\omega_n^{(k)}, \omega_m^{(l)}] = \sum_{p \geq 0} A_{kl}^p \omega_{m+n}^{(k+l-2p-1)},$$

where $A_{kl}^p = \sum_{i+j=2p+1} (-1)^i C_{k-1}^i C_{l-1}^j$, $C_n^m = \frac{n!}{m!(n-m)!}$ if $m \leq n$ and 0 otherwise.

Now let us mention that all our constructions admit certain limits with one or both parameters N and M tends to infinity. That means that the torus becomes a cylinder or a plane. In the case $M \rightarrow \infty$ the Poisson algebra $\mathcal{F}^{\mathbb{C}}(\mathbb{T}^2)$ inverts into the Poisson algebra $\mathcal{F}^{\mathbb{C}}(T^*\mathbb{S}^1)$ of functions on a cylinder. There is a natural basis in $\mathcal{F}^{\mathbb{C}}(T^*\mathbb{S}^1)$: $e_n^m = -i \exp(\frac{inq}{N}) p^m$, in which the Poisson brackets have the form:

$$\{e_n^m, e_{n'}^{m'}\} = (mn' - nm') e_{n+n'}^{m+m'-1}.$$

The embedding of the algebra $\mathbb{C}\text{Vect}(\mathbb{S}^1)$ into $\mathcal{F}^{\mathbb{C}}(T^*\mathbb{S}^1)$ is natural: $L_k \mapsto e_k^1$. Such embedding admits, indeed an one-parametric deformation $L_k \mapsto e_k^1 + (k+1)\lambda e_k^0$, λ is a parameter. The central extension $\widehat{\mathcal{F}}^{\mathbb{C}}(T^*\mathbb{S}^1)$ of $\mathcal{F}^{\mathbb{C}}(T^*\mathbb{S}^1)$ has the form:

$$\{e_n^m, e_{n'}^{m'}\} = (mn' - nm') e_{n+n'}^{m+m'-1} + \frac{1}{12} (n^3 - n) \delta(n+n') \delta_{m,1} \delta_{m',1} \mathbf{1}$$

The embeddings of $\mathbb{C}\text{Vect}(\mathbb{S}^1)$ into $\mathcal{F}^{\mathbb{C}}(T^*\mathbb{S}^1)$ extends to embeddings of the Virasoro algebra $\mathbb{C}\text{vir}$ into $\widehat{\mathcal{F}}^{\mathbb{C}}(T^*\mathbb{S}^1)$.

The Poisson algebra $\mathcal{F}^{\mathbb{C}}(T^*\mathbb{S}^1)$ is deformed into the algebra $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)$ of differential operators on a circle. The corresponding commutator algebra is a Lie algebra $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ of differential operators on a circle. The embeddings of $\mathbb{C}\text{Vect}(\mathbb{S}^1)$ in $\mathcal{F}^{\mathbb{C}}(T^*\mathbb{S}^1)$ induce embeddings of this algebra into $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$; these embedding are the same if one consider a subalgebra $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$ of $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$, the commutator algebra of the algebra $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_+$ of differential operators on the circle $\mathbb{S}^1$ without free terms. Both algebras $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ and $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$ admit central extensions $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ and $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$, which are defined by the Kac-Peterson cocycle $c(f(q)D^m, g(q)D^n) = \frac{m!n!}{(m+n+1)!} \int_{\mathbb{S}^1} f^{(n)}(q)g^{(m+1)}(q) dq$, where $D = \partial/\partial q$, $f^{(n)}(q) = \frac{\partial^n f(q)}{\partial q^n}$ [21] (see also [6]). The embeddings of $\mathbb{C}\text{Vect}(\mathbb{S}^1)$ in $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ (resp. the emdedding in $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$) are extended to embeddings of the Virasoro algebra $\mathbb{C}\text{vir}$ in $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ (resp. an embedding in $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$).

To make the construction of central extensions $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ and $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$ of $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ and $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$ more clear one should transit to the algebra $\text{PDOP}^{\mathbb{C}}(\mathbb{S}^1)$ of all pseudodifferential operators on a circle $\mathbb{S}^1$ [5-8]. The commutator algebra $\text{PDOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ of this algebra, the Lie algebra of pseudodifferential operators on a circle, admits a central extension, which may be defined as $[A, B]_c = [A, B] + c \cdot \text{Tr}([A, \log D], B)$, where $\text{Tr}(A) = \int \text{Res}(A) dq$, $\text{Res}(A) = a_{-1}(q)$ if $A = \sum_{k \in \mathbb{Z}} a_k(q) D^k$ [5]. The cocycle $c(A, B) = \text{Tr}([A, \log D], B)$ is so-called

Kravchenko-Khesin cocycle. The extension by Kravchenko-Khesin cocycle being restricted on subalgebras $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ and $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$ define extensions described above.

In the case $M, N \rightarrow \infty$ the Poisson algebra $\mathcal{F}^{\mathbb{C}}(\mathbb{T}^2)$ inverts into the Poisson algebra $\mathcal{F}^{\mathbb{C}}(\mathbb{R}^2)$ of functions on a plane $\mathbb{R}^2$. There is defined a natural polynomial basis in $\mathcal{F}^{\mathbb{C}}(\mathbb{R}^2)$: $e^{mn} = p^m q^n$, in which the Poisson brackets have the form

$$\{e^{mn}, e^{m'n'}\} = (mn' - nm')e^{m+m'-1, n+n'-1}.$$

The Poisson algebra $\mathcal{F}^{\mathbb{C}}(\mathbb{R}^2)$ may be deformed into the Weyl algebra $W_{\mathbb{C}}(\mathbb{R}^2)$. Polynomial elements of this algebra may be represented by differential operators on a line $\mathbb{R}^1$, so there is corresponded a certain differential operator $\mathcal{D}_P$ to the polynomial P from $\mathcal{F}^{\mathbb{C}}(\mathbb{R}^2)$, where P is a Weyl (symmetric) symbol of $\mathcal{D}_P$. The commutator algebra of the Weyl algebra $W_{\mathbb{C}}(\mathbb{R}^2)$ is so-called Moyal algebra $\text{Moy}_{\mathbb{C}}(\mathbb{R}^2)$. The commutator in the Moyal algebra has the form [19,22]

$$[F, G] = \sum_{p \in \mathbb{Z}} (-1)^p \sum_{i+j=2p+1} \frac{(-1)^i}{i! j!} (\partial_p^i \partial_q^j F) (\partial_p^j \partial^i G)$$

or in the basis e^{mn}

$$[e^{mn}, e^{m'n'}] = \sum_{p \geq 0} (-1)^p A_{mm'nn'}^{2p+1} e^{m+m'-2p-1, n+n'-2p-1},$$

where $A_{mm'nn'}^p = \sum_{i+j=p} (-1)^i i! j! C_n^i C_n^j C_{m'}^j C_{n'}^i$, $C_n^m = \frac{n!}{m!(n-m)!}$ if $m \leq n$ and 0 otherwise.

1.2. Poisson algebra $\mathcal{F}^{\mathbb{C}}(T^*\mathbb{S}^1)$, its central extension $\widehat{\mathcal{F}}^{\mathbb{C}}(T^*\mathbb{S}^1)$ and w_{∞} -algebra (Bakas algebra); Lie algebra $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$, its central extension $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ and W_{∞} -algebras (Pope-Shen-Romans algebras). Gelfand-Dickey algebras $\text{GD}_n^{\mathbb{C}}$ and Radul mapping $\mathcal{F}(\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)^*) \mapsto \text{GD}_n^{\mathbb{C}}$; Radul bundle $\text{Rad}^{\mathbb{C}}(\mathcal{M}_n)$ and the Lie algebra of its sections $\text{Rad}_{\mathbb{C}}$ (Radul algebra). Wedge subalgebra of Pope-Sheen-Romans W_{∞} -algebra; associative model algebras $\text{Md}^{(\lambda)}(\text{sl}(2, \mathbb{C}))$ for the Lie algebra $\text{sl}(2, \mathbb{C})$ and their commutator algebras $\text{Md}_{[\cdot, \cdot]}^{(\lambda)}(\text{sl}(2, \mathbb{C}))$ - model Lie algebras (or Feigin algebras) for $\text{sl}(2, \mathbb{C})$; Racah-Wigner algebras $\mathcal{RW}_{\infty}^{(\lambda)}(\text{sl}(2, \mathbb{C}))$ for $\text{sl}(2, \mathbb{C})$, their reductions $\mathcal{RW}_n^{(\lambda)}(\text{sl}(2, \mathbb{C}))$ of order n and central extensions $\widehat{\mathcal{RW}}_n^{(\lambda)}(\text{sl}(2, \mathbb{C}))$ of $\mathcal{RW}_n^{(\lambda)}(\text{sl}(2, \mathbb{C}))$.

The Poisson algebra $\mathcal{F}^{\mathbb{C}}(\mathbb{S}^1)$ may be expanded under the action of $\mathbb{C}\text{Vect}(\mathbb{S}^1)$ into the sum of tensor modules of positive integer weights, the corresponding tensor operators $w_k^{(i)}$ ($i \geq 1$) are equal to e_k^{i-1} so that the commutation relations

$$[w_m^{(i)}, w_n^{(j)}] = ((j-1)n - (i-1)m)w_{n+m}^{(i+j-2)}$$

holds. The algebra spanned by $w_n^{(i)}$, $n \in \mathbb{Z}$, $i \geq 2$ is closed and is called w_{∞} -algebra (or *Bakas algebra*) [23] as well as its central extension, which is obtained from $\widehat{\mathcal{F}}^{\mathbb{C}}(T^*\mathbb{S}^1)$.

Now we are interested in the deformations of w_{∞} -algebra, when $\widehat{\mathcal{F}}^{\mathbb{C}}(T^*\mathbb{S}^1)$ deforms into $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$. Such deformations were found by C.N.Pope, X.Shen and L.J.Romans [24] (see also [25]), so the corresponding algebras are called *Pope-Shen-Romans-algebras* (or W_{∞} -algebras). The explicit formulas for their generators were found by I.Bakas, B.Khesin and E.Kiritsis [8] so the embedding of W_{∞} -algebras in $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ are called *Bakas-Khesin-Kiritsis embeddings*.

Let us at first construct so-called $W_{1+\infty}$ -algebra, which is an algebra of $\mathbb{C}\text{vir}$ -tensor operators in $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$, following [8]. Its generators V_m^s are labelled by $m \in \mathbb{Z}$ and $s \geq 1$. Their explicit form is

$$V_m^s = -B(s) \sum_{k=1}^s \alpha_k^s C_{k-1}^{m+s-1} z^{m+s-k} D_z^{s-k},$$

where $B(s) = \frac{2^{s-3}(s-1)!}{(2s-3)!!}$, $\alpha_k^s = \frac{(2s-k-1)!}{[(s-k)!]^2}$; $z = \exp(iq)$, $D_z = \frac{\partial}{\partial z}$. To describe commutation relations in $W_{1+\infty}$ -algebra it is convenient to introduce some notations

$$\begin{aligned} g_l^{ss'}(m, n; \mu) &:= \frac{1}{2(l+1)!} f_l^{ss'}(\mu) N_l^{ss'}(m, n) \\ f_l^{ss'}(\mu) &:= \sum_{k \geq 0} \frac{(-\frac{1}{2} - 2\mu)_k (\frac{3}{2} + 2\mu)_k (-\frac{l+1}{2})_k (-\frac{l}{2})_k}{k! (-s + \frac{3}{2})_k (-s' + \frac{3}{2})_k (s+s'-l-\frac{3}{2})_k} \\ N_l^{ss'}(m, n) &:= \sum_{k=0}^{l+1} (-1)^k C_k^{l+1} (2s-l-2)_k [2s'-k-2]_{l+1-k} [s-1+m]_{l+1-k} [s'-1+n]_k \end{aligned}$$

where $(a)_k := a(a+1) \dots (a+k-1)$, $[a]_k := a(a-1) \dots (a-k+1)$. Then

$$\begin{aligned} [V_m^s, V_n^{s'}] &= ((s'-1)m - (s-1)n) V_{m+n}^{s+s'-2} + c_s(m, \mu) \delta_{ss'} \delta(m+n) + \\ &\quad \sum_{r \geq 0} g_{2r}^{ss'}(m, n; \mu) V_{m+n}^{s+s'-2r-2}, \end{aligned}$$

where $\mu = -\frac{1}{2}$,

$$c_s(m; -\frac{1}{2}) = c \cdot \frac{(m+s-1)!}{(m-s)!} \cdot \frac{2^{2(s-3)}[(s-1)!]^2}{(2s-1)!!(2s-3)!!}.$$

$W_{1+\infty}$ -algebra is not a deformation of Bakas algebra because it contains tensor operators of upper index 1. To construct the correct deformation (which is the "right" W_∞ -algebra) one should transform generators V_m^s in the following manner. Let us introduce new generators W_m^s ($m \in \mathbb{Z}$, $s \geq 2$) by the formulas

$$W_m^s = V_m^s + \frac{B(s)}{s-1} \sum_{k=1}^{s-1} (-1)^l \frac{(2s-2l-1)}{B(s-l)} \cdot \frac{(m+s-1)!}{(m+s-l-1)!} V_m^{s-1}$$

or explicitly

$$W_m^s = -\frac{B(s)}{s-1} \sum_{k=1}^{s-1} \beta_k^s C_{k-1}^{m+s-1} z^{m+s-k} D_z^{s-k},$$

where $\beta_k^s = \frac{(2s-k-1)!}{(s-k)!(s-k-1)!}$, i.e. W_∞ -algebra is realized in $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$. The commutation relations in W_∞ -algebra have the form

$$[W_m^s, W_n^{s'}] = ((s'-1)m - (s-1)n) W_{m+n}^{s+s'-2} + c_s(m; \mu) \delta_{ss'} \delta(m+n) + \sum_{r \geq 1} g_{2r}^{ss'}(m, n; \mu) W_{m+n}^{s+s'-2-2r},$$

where $\mu = 0$,

$$c_s(m, 0) = \frac{c}{2} \cdot \frac{(m+s-1)!}{(m-s)!} \cdot \frac{2^{2(s-3)} s! (s-2)!}{(2s-1)!! (2s-3)!!}.$$

Let us consider following [6,7] an affine subspace $\mathcal{M}_n$ in the algebra $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)$ of the form $\mathcal{M}_n = \{A \in \text{DOP}^{\mathbb{C}}(\mathbb{S}^1) : A = D^n + a_{n-1}(q)D^{n-1} + \dots + a_1(q)D + a_0(q)\}$. The tangent space $T_A(\mathcal{M}_n)$ of manifold $\mathcal{M}_n$ at the point A may be identified with the space of all operators X from $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)$ of the form $X = x_{n-1}(q)D^{n-1} + \dots + x_0(q)$. The corresponding cotangent space $T_A^*(\mathcal{M}_n)$ may be identified with the space of all operators Y from $\text{PDOP}^{\mathbb{C}}(\mathbb{S}^1)$ of the form $Y = y_{-1}(q)D^{-1} + \dots + y_{1-n}(q)D^{1-n} + y_{-n}(q)D^{-n}$. The spaces $T_A^*(\mathcal{M}_n)$ and $T_A(\mathcal{M}_n)$ are paired by $\langle X, Y \rangle = \text{Tr}(XY)$.

There is defined [6,7] a tensor operator field $V : T^*(\mathcal{M}_n) \mapsto T(\mathcal{M}_n)$, namely V_A , which maps $T_A^*(\mathcal{M}_n)$ to $T_A(\mathcal{M}_n)$, is defined by the following formulas $V_A(Y) = A(YA)_+ - (AY)_+A$, $Y \in T_A^*(\mathcal{M}_n)$, where $A \mapsto A_+$ is the natural projection of $\text{PDOP}^{\mathbb{C}}(\mathbb{S}^1)$ to $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)$.

The space $\mathcal{F}(\mathcal{M}_n)$ of functions on the manifold $\mathcal{M}_n$ possesses a structure of the Poisson algebra via $\{F, G\} = \langle V(dF), dG \rangle$ [6,7]. This Poisson algebra is called *Gelfand-Dickey algebra* and is denoted by $\text{GD}_n^{\mathbb{C}}$. The center of Gelfand-Dickey algebra $\text{GD}_n^{\mathbb{C}}$ consists of constant functions, so the strict sequence

$$0 \longrightarrow \mathbb{C} \longrightarrow \text{GD}_n^{\mathbb{C}} \longrightarrow \overline{\text{GD}}_n^{\mathbb{C}} \longrightarrow 0$$

exists, where $\overline{\text{GD}}_n^{\mathbb{C}}$ is a quotient $\text{GD}_n^{\mathbb{C}}/\mathbb{C}$.

Let us now construct the *Radul mapping* from the Poisson algebra $\mathcal{F}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)^*)$ of the functions on the coadjoint module $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)^*$ of the Lie algebra $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ onto the Poisson algebra $\text{GD}_n^{\mathbb{C}}$. Namely, let us construct a mapping $H : \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1) \mapsto \mathbb{C}\text{Vect}(\mathcal{M}_n)$, where the mapping $H_A : \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1) \mapsto T_A(\mathcal{M}_n)$ is defined by the formulas $H_A(B) = V_{(BA^{-1})_-}(A) = AB - (ABA^{-1})_+A$; $B \in \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)$, $A_- = A - A_+$ [6,7]. It should be mentioned that $\text{Ker}(H_A) = \{C \in \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1) : C = BA, A \in \text{DOP}^{\mathbb{C}}(\mathbb{S}^1)\}$. It is stated that $\text{Im}(H)$ is contained in $\text{Im}(d)$, where $d : \text{GD}_n^{\mathbb{C}} \mapsto \mathbb{C}\text{Vect}(\mathcal{M}_n)$. Because $\text{Ker}(d)$ consists of constant functions on $\mathcal{M}_n$ then H realizes a homomorphism of $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ into $\overline{\text{GD}}_n^{\mathbb{C}}$ considered as a Lie algebra. Such homomorphism is extended to the homomorphism of the Poisson algebra $\mathcal{F}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)^*)$ onto the Gelfand-Dickey algebra $\text{GD}_n^{\mathbb{C}}$. Of course, one may consider a restriction of the Radul mapping $\mathcal{F}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)^*) \mapsto \text{GD}_n^{\mathbb{C}}$ on the Poisson subalgebra $\mathcal{F}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^*)$ of the Poisson algebra $\mathcal{F}(\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)^*)$, where $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_+$ is a subalgebra of $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)$ of differential operators without free terms, $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$ is its commutator algebra, the Lie algebra of differential operators without free terms, and $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^*$ is the coadjoint module over this Lie algebra.

Now let us construct so-called *Radul bundle* and the Lie algebra of its sections (*Radul algebra*). Let us consider a trivial bundle $\text{Rad}^{\mathbb{C}}(\mathcal{M}_n)$ over $\mathcal{M}_n$ with fiber isomorphic to $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)$. One may define the following commutator in the space of its sections: if $E, F \in \mathcal{O}(\text{Rad}^{\mathbb{C}}(\mathcal{M}_n))$ then

$$[E, F]'(A) := [E(A), F(A)] + V_{(EA^{-1})_-}(F) - V_{(FA^{-1})_-}(E) \in \text{Rad}_A^{\mathbb{C}}(\mathcal{M}_n),$$

$$\text{where } V_X(E(A)) = \left. \frac{\partial}{\partial t} E(A + tV_X(A)) \right|_{t=0}, \quad A \in \mathcal{M}_n.$$

The algebra of sections of the bundle $\text{Rad}^{\mathbb{C}}(\mathcal{M}_n)$ will be called the Radul algebra $\text{Rad}_{\mathbb{C}}$. As it was stated in [7] the mappings H_A being glued define a homomorphism of the Radul algebra $\text{Rad}_{\mathbb{C}}$ onto $\overline{\text{GD}}_n^{\mathbb{C}}$ considered as a Lie algebra.

The algebra $\text{GD}_n^{\mathbb{C}}$ defined above is just Gelfand-Dickey algebra $\text{GD}(\text{gl}(n, \mathbb{C}))$ for the Lie algebra $\text{gl}(n, \mathbb{C})$. It is more convenient to consider Gelfand-Dickey algebra $\text{GD}(\text{sl}(n, \mathbb{C}))$ for the Lie algebra $\text{sl}(n, \mathbb{C})$. To do it one should consider the subspace $\mathcal{M}_n^{(0)}$ of $\mathcal{M}_n$ of operators A with $a_{n-1} = 0$, it should be mentioned that our presentation may be considered for $\text{GD}(\text{sl}(n, \mathbb{C}))$ if one change $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ on $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$.

As it was marked in [24] generators W_m^s of the Pope-Shen-Romans W_{∞} -algebra with $1 - s \leq m \leq s - 1$ form a closed Lie algebra, which is called *wedge subalgebra* of W_{∞} -algebra. The generators W_{-1}^2, W_0^2, W_1^2 of the wedge subalgebra are just the generators of the Lie algebra $\text{sl}(2, \mathbb{C})$. A space of the wedge subalgebra as $\text{sl}(2, \mathbb{C})$ -module is identified with the model $M_{\text{odd dim}}(\text{sl}(2, \mathbb{C}))$ of odd-dimensional representations of the Lie algebra $\text{sl}(2, \mathbb{C})$ [26].

Let us analyse the algebraic structures related to the model $M_{\text{odd dim}}(\text{sl}(2, \mathbb{C}))$ more systematically. If one considered the universal enveloping algebra $\mathcal{U}(\text{sl}(2, \mathbb{C}))$ of the Lie algebra $\text{sl}(2, \mathbb{C})$ and the ideal I_{λ} , generated by $K - \lambda$, where K is a Casimir

operator for $\mathfrak{sl}(2, \mathbb{C})$ (i.e. a quadratic element of the center $\mathcal{Z}(\mathcal{U}(\mathfrak{sl}(2, \mathbb{C})))$ of the universal enveloping algebra $\mathcal{U}(\mathfrak{sl}(2, \mathbb{C}))$), in it, then the quotient $\mathcal{U}(\mathfrak{sl}(2, \mathbb{C}))/I_\lambda$ supplies the model $M_{\text{odd dim}}(\mathfrak{sl}(2, \mathbb{C}))$ by a structure of an associative algebra. Such associative algebra will be called *the associative model algebra for the Lie algebra* $\mathfrak{sl}(2, \mathbb{C})$ and will be denoted by $\text{Md}^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$.

The commutator algebra $\text{Md}_{[\cdot, \cdot]}^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ of the associative model algebra $\text{Md}^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ will be called *the model Lie algebra for the Lie algebra* $\mathfrak{sl}(2, \mathbb{C})$ (or *Feigin algebra*, see [27]). The wedge subalgebra of the Pope-Shen-Romans W_∞ -algebra is just the Feigin algebra $\text{Md}_{[\cdot, \cdot]}^{(0)}(\mathfrak{sl}(2, \mathbb{C}))$.

Let us now consider the Racah-Wigner algebras, their reductions of finite order and central extensions of such reductions for the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$. *The Racah-Wigner algebra* $\mathcal{RW}_\infty^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ *for the Lie algebra* $\mathfrak{sl}(2, \mathbb{C})$ is the universal enveloping algebra $\mathcal{U}(\text{Md}_{[\cdot, \cdot]}^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C})))$ of the model Lie algebra $\text{Md}_{[\cdot, \cdot]}^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$.

Definition 1. An associative algebra $\mathcal{RW}_n^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ is called *a reduced Racah-Wigner algebra of order n for the Lie algebra* $\mathfrak{sl}(2, \mathbb{C})$ iff (1) it admits a homomorphism onto $\text{Md}^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$; (2) $\mathcal{RW}_n^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ is generated by the direct sum $\pi_1 \oplus \pi_2 \oplus \dots \oplus \pi_{n-1} \oplus \pi_n$ of the first n odd-dimensional representations of $\mathfrak{sl}(2, \mathbb{C})$ (this direct sum is isomorphic to $\mathfrak{sl}(n, \mathbb{C})$ as $\mathfrak{sl}(2, \mathbb{C})$ -module), the natural action of $\mathfrak{sl}(2, \mathbb{C})$ is defined by $\text{ad}(\pi_1)$; (3) $\mathcal{RW}_n^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ as $\mathfrak{sl}(2, \mathbb{C})$ -module is isomorphic to $S(\mathfrak{sl}(n, \mathbb{C}))$ ($S(V)$ is the symmetric algebra over V).

Reduced Racah-Wigner algebras $\mathcal{RW}_2^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ of order 2 with non-homogeneous quadratic relations were considered in [28]. It was shown that $\mathcal{RW}_2^{(\lambda_1)}(\mathfrak{sl}(2, \mathbb{C})) \simeq \mathcal{RW}_2^{(\lambda_2)}(\mathfrak{sl}(2, \mathbb{C}))$ for arbitrary λ_1 and λ_2 .

Proposition 1. *There exists a reduced Racah-Wigner algebra $\mathcal{RW}_n^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ of finite order n for the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ for all n .*

Proof. Unfortunately, we do not know the direct algebraic proof of this proposition, which is, certainly, preferable. Let us formulate a quantum-field one, which is based on formalism of the q_R -conformal field theory [28]. Namely, let us consider the set of q_R -affine currents, which charges form the Lie algebra $\mathfrak{sl}(n, \mathbb{C})$ (or which components form q_R -affine Lie algebra) [28]. Let us consider the set of Casimir operators $K_1, \dots, K_{n-1}$ and the higher spin fields $W^2(z), \dots, W^n(z)$, which correspond to them in the operator algebra of q_R -affine currents. The algebra, generated by the components W_m^s ($1 - s \leq m \leq s - 1$; $W^s(z) = \sum_{m \in \mathbb{Z}} W_m^s z^{-s-m}$) is just the reduced Racah-Wigner algebra $\mathcal{RW}_n^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ of the order n for the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$, where $\lambda = \frac{1}{4}(q_R^{-1} + 3)(q_R^{-1} + 1)$.

The non-linear structure constants of the constructed reduced Racah-Wigner algebras $\mathcal{RW}_n^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ admits a deformation by the linear structure constants of the universal enveloping algebra $\mathcal{U}(\mathfrak{sl}(n, \mathbb{C}))$ of the Lie algebra $\mathfrak{sl}(n, \mathbb{C})$. One may introduce a new central element ρ and consider this deformation as a central extension of the reduced Racah-Wigner algebra. Such central extension will be denoted by $\widehat{\mathcal{RW}}_n^{(\lambda)}(\mathfrak{sl}(n, \mathbb{C}))$. If one consider the fixed value of central element ρ , i.e. a quotient of $\widehat{\mathcal{RW}}_n^{(\lambda)}(\mathfrak{sl}(n, \mathbb{C}))$ by the ideal generated by $\rho - \rho_0$, then the limit $\rho_0 \rightarrow \infty$ inverts the obtained quotient into the universal enveloping algebra

$\mathcal{U}(\mathfrak{sl}(n, \mathbb{C}))$. So one may consider the algebras $\widehat{\mathcal{RW}}_n^{(\lambda)}(\mathfrak{sl}(2, \mathbb{C}))$ as certain deformations of $\mathcal{U}(\mathfrak{sl}(n, \mathbb{C}))$.

2. GÉRAIS-MATSUO DIFFERENTIAL W -GEOMETRY AND W -SYMMETRIES OF A SECOND QUANTIZED FREE STRING

In this paragraph we shall work with the following objects [1](see also [13,29-31]):

- Q (or the dual Q^*) — the space of external degrees of a freedom of a string. The coordinates x_n^μ on Q are the Taylor coefficients of functions $x^\mu(z)$, which determines a world-sheet of a string in a complexified target space.
- $M(\text{Vir})$ — the space of internal degrees of a freedom of a string; *the flag manifold of the Virasoro-Bott group* Vir ; the homogeneous space $\text{Diff}_+(\mathbb{S}^1)/\mathbb{S}^1$ ($\text{Diff}_+(\mathbb{S}^1)$ is the group of diffeomorphisms of a circle $\mathbb{S}^1$ preserving an orientation); this space is identified via Kirillov construction [13;A.A.Kirillov] with
- S — *the class of the univalent functions* $f(z)$ in the unit complex disc D_+ ($D_+ = \{z \in \mathbb{C}: |z| \leq 1\}$) such that $f(0) = 0$, $f'(0) = 1$, $f'(e^{it}) \neq 0$; the natural coordinates on S are coefficients c_k of the Taylor expansion of an univalent function $f(z)$: $f(z) = z + c_1 z^2 + c_2 z^3 + \dots + c_{n-1} z^n + c_n z^{n+1} + \dots$
- $\mathcal{C}$ — *the universal deformation of a complex disc* with $M(\text{Vir})$ as a base and with fibers isomorphic to D_+ . The coordinates on $\mathcal{C}$ are $z, c_1, c_2, \dots, c_n, \dots$, where c_k are coordinates on the base and z is a coordinate in the fibers.
- $M(\text{Vir}) \cdot Q^*$ — the space of both external and internal degrees of freedom of a string, the same as the bundle over $M(\text{Vir})$ associated with $p: \mathcal{C} \mapsto M(\text{Vir})$, which fibers are $\text{Map}(\mathcal{C}/M(\text{Vir}); \mathbb{C}^n)^*$ — linear spaces dual to ones of mappings of fibers of $p: \mathcal{C} \mapsto M(\text{Vir})$ into $\mathbb{C}^n$.

2.1. Elements of Gervais-Matsuo differential W -geometry: classical Toda fields in the complex analogue of Frenet theory; the mapping $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}} \mapsto \mathbb{C}\text{Vect}(M(\text{Vir}) \cdot Q^*)$; Gervais-Matsuo Lie quasi(pseudo)algebra $\mathcal{GM}_n^{\mathbb{C}}$ of classical (infinitesimal) W -transformations; Gervais-Matsuo Poisson algebra $\text{GM}_n^{\mathbb{C}}$ and the monomorphism $\text{GM}_n^{\mathbb{C}} \mapsto \text{GD}(\mathfrak{sl}(n, \mathbb{C}))$. Infinite dimensional geometry of the flag manifold $M(\widehat{\text{DOP}}_{[\cdot, \cdot]}(\mathbb{S}^1)_+) \simeq M(W_\infty^r)$ ($W_\infty = (W_\infty^r)^{\mathbb{C}}$) for the Lie algebra $\widehat{\text{DOP}}_{[\cdot, \cdot]}(\mathbb{S}^1)_+$ or for the real form W_∞^r of the Pope-Shen-Romans W_∞ -algebra.

Let us consider a world-sheet of a string $\mathbf{x}: D_+ \mapsto \mathbb{C}^n$ ($\mathbf{x} = \{x^\mu = x^\mu(z)\}$), then using the complex version of Frenet theory [12] one may introduce the associate mappings $\text{Gr}_k \mathbf{x}: D_+ \mapsto \text{Gr}_k(\mathbb{C}^n)$, where $\text{Gr}_k(\mathbb{C}^n)$ is the Grassmannian of k -dimensional planes in $\mathbb{C}^n$, or the associated mapping $\text{Fl} \mathbf{x}: D_+ \mapsto \text{Fl}(\mathbb{C}^n)$, where $\text{Fl}(\mathbb{C}^n)$ is the space of full flags in $\mathbb{C}^n$. If we consider the homogeneous coordinates or go to the projective space $\mathbb{CP}^n$, then *the classical Toda fields* on D_+ will be identified with Kähler potentials of the images of $\text{Gr}_{k+1} \mathbf{x}(z)$ in $\text{Gr}_{k+1}(\mathbb{C}^{n+1})$.

It is convenient to introduce so-called *homogeneous KP-coordinates* for the mapping $\mathbf{x}: D_+ \mapsto \mathbb{CP}^n$. Indeed, let us consider the functions $x^\mu([z])$, $[z] = [z^{(0)}, z^{(1)} = z, z^{(2)}, \dots, z^{(n)}, \dots]$ such that $D_z^l \mathbf{x}([z]) = \frac{\partial \mathbf{x}([z])}{\partial z^{(l)}}$ and $\mathbf{x}([z]) = \mathbf{x}(z)$ for $z^{(2)}, z^{(3)}, \dots, z^{(n)}, \dots = 0, z = z^{(1)}/z^{(0)}$. So the homogeneous KP-coordinates $[z^{(0)}, z^{(1)}, \dots, z^{(n)}]$ may be regarded as certain ones in the complex projective space $\mathbb{CP}^n$ defined in the neighbourhood of the world-sheet of a string.

Let now $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)^{\text{reg}}$ be a subalgebra of $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)$, which consists of operators possessing a regular continuation to the unit complex disc D_+ ; $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}$ be the intersection of $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)^{\text{reg}}$ with $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_+$; $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)^{\text{reg}}$ and $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}$ be the corresponding commutator Lie algebras.

It should be mentioned that $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}$ naturally acts by differential operators on the space Q^* , considered as the space of sections of a trivialized bundle over D_+ with fibers isomorphic to $\mathbb{C}^n$. One of the results of J.-L.Gervais and Y.Matsuo was that this action is linearized (i.e. becomes an action by vector fields) in KP-coordinates.

Proposition 2. *The action of the Lie algebra $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}$ by differential operators on Q^* (as the space of sections of a trivialized bundle over D_+ with fibers isomorphic to $\mathbb{C}^n$) may be extended to the action on $M(\text{Vir}) \cdot Q^*$ (as the space of sections of a trivialized bundle over $\mathcal{C}$ with fibers isomorphic to $\mathbb{C}^n$) by vector fields.*

Proof. It is necessary just to mention that KP-coordinate system is a homogeneous version of the coordinates on the universal deformation $\mathcal{C}$ of a complex disc D_+ , which are easily expressed via the standard coordinates $z, c_1, c_2, \dots, c_n, \dots$.

Let us now consider more systematically the structure of the action of the Lie algebra $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}$ on Q^* by differential operators.

It is rather reasonable following J.-L.Gervais and Y.Matsuo to restrict ourselves to a consideration of differential operators of order less or equal to n . It means that we shall deal with the quotient $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}} / \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}}$ of the Lie algebra $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}$ of regular differential operators without free terms by its subalgebra $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}}$ of regular differential operators without free terms, which are not contained terms with D^k ($1 \leq k \leq n$), the commutator algebra of associative algebra $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}}$ of such operators. Of course, such quotient is not a Lie algebra, because the subalgebra $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}}$ is not an ideal in $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}$. Nevertheless, the strict sequence

$$0 \longrightarrow \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}} \longrightarrow \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}} \longrightarrow \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}} / \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}} \longrightarrow 0$$

may be resolved. The resolving mapping identifies the quotient $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}} / \text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}}$ with the subspace $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_{+;\leq n}^{\text{reg}}$ of regular differential operators of order less or equal to n without free terms in the Lie algebra $\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}$.

The object, which elements of $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_{+;\leq n}^{\text{reg}}$ form is, indeed, a Lie quasialgebra (in terminology of I.Batalin [32] or Lie pseudoalgebra in terminology of M.V.Karasev and V.P.Maslov [33]). Namely, the commutator of two elements of $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_{+;\leq n}^{\text{reg}}$ acting on Q^* (classical (infinitesimal) W -transformations [11]) may be expressed via other elements with coefficients, which are extrinsic invariants of the world-sheet of a string (curvature, torsions and their derivatives). So the structure functions of classical (infinitesimal) W -transformations are the functions on the space Q^* , the object, on which classical (infinitesimal) W -transformations act. Therefore, classical (infinitesimal) W -transformations form a Lie quasi(pseudo)algebra, which will be called *Gervais-Matsuo quasi(pseudo)algebra* and will be denoted by $\mathcal{GM}_n^{\mathbb{C}}$.

It should be marked (with respect to the possible constructing of classical finite W -transformations) that Lie quasi(pseudo)algebras are infinitesimal objects for Lie quasi(pseudo)groups of transformations in the finite dimensional case [32,33]. Fixing a point of a manifold, on which the Lie quasi(pseudo)group of transformations acts, one may receive a structure of a loop on the space of the Lie quasi(pseudo)-group. In our case this loop will be just the loop of homogeneous space $\text{EXP}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}})/\text{EXP}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}})$ defined by $\text{EXP}(\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_{+;\leq n}^{\text{reg}})$ as a resolution of the strict sequence

$$0 \longrightarrow \text{EXP}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}}) \longrightarrow \text{EXP}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}) \longrightarrow \\ \text{EXP}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}})/\text{EXP}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}}) \longrightarrow 0$$

via *Sabinin construction* [34]. Nevertheless, infinite dimensional groups $\text{EXP}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}^{\text{reg}})$ and $\text{EXP}(\text{DOP}_{[\cdot,\cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}})$ (as I know) are not constructed and, therefore, the classical finite W -transformations are not introduced; so Sabinin construction may be considered only as an explanation of the appearance of Gervais-Matsuo quasi(pseudo)algebra. However, it should be marked, though the Lie quasi(pseudo)group of classical finite W -transformations is not defined (though the author thinks that it is possible) and, hence, the corresponding system of isotopic loops on it can not be considered, the infinitesimal objects of such loops, so called *Mikheev-Sabinin multialgebras* [35] may be derived only from Gervais-Matsuo quasi(pseudo)algebra $\mathcal{GM}_n^{\mathbb{C}}$.

It should be mentioned that Q^* is a symplectic manifold and, therefore, one may correspond to the generators of Gervais-Matsuo quasi(pseudo)algebra $\mathcal{GM}_n^{\mathbb{C}}$ their Hamiltonians, which are just functions of classical Toda fields. The structure of the Lie quasi(pseudo)algebra induces a structure of the Poisson algebra on Hamiltonians, which will be called *Gervais-Matsuo Poisson algebra* and will be denoted by $\text{GM}_n^{\mathbb{C}}$. There exists a very simple but remarkable fact, which we prefer to formulate as a proposition.

Proposition 3. *There exist a monomorphism of Poisson algebras*

$$\text{GM}_n^{\mathbb{C}} \mapsto \text{GD}(\mathfrak{sl}(n, \mathbb{C}))$$

Namely, Gervais-Matsuo algebra $\text{GM}_n^{\mathbb{C}}$ is just a "regular" part of Gelfand-Dickey algebra $\text{GD}(\mathfrak{sl}(n, \mathbb{C}))$.

The significance of such monomorphism is explained by the fact that *Gervais-Matsuo differential W -geometry may be (at least, partially) generalized on an arbitrary Kähler manifold as a complexified target space*, because it is based essentially only on complex Frenet theory (see [11,36]), so one can construct analogs of Gervais-Matsuo quasi(pseudo)algebra $\mathcal{GM}_n^{\mathbb{C}}$ of classical (infinitesimal) W -transformations as well as Gervais-Matsuo Poisson algebra $\text{GM}_n^{\mathbb{C}}$ for an arbitrary Kähler manifold.

It is not less remarkable circumstance that *Gervais-Matsuo differential W -geometry appeared being deeply connected with Sabinin program of "nonlinear geometric algebra" [37] and Weinstein-Karasev-Maslov approach to nonlinear Poisson brackets [33]; so from the point of view of algebraic geometry Gervais-Matsuo W -geometry may be considered as a penetration of nonassociative algebra into the theory of embeddings of algebraic curves into Kähler varieties* (it is reminiscent a little of a book of Yu.I.Manin [38]).

But let us now return to the main theme. Above we obtained that the subalgebra W_∞^{reg} of W_∞ -algebra, generated by V_m^s , $m \geq 0$ acts by vector fields on the space $M(\text{Vir}) \cdot Q^*$. Nevertheless, we want to obtain an action of the whole W_∞ -algebra (or $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$) instead of W_∞^{reg} (or $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+^{\text{reg}}$) only. So we should to enlarge the space of the internal symmetries of a string by new degrees of freedom. What we shall to do is to consider the flag manifold $M(\widehat{\text{DOP}}_{[\cdot, \cdot]}(\mathbb{S}^1)_+)$ for the Lie algebra $\widehat{\text{DOP}}_{[\cdot, \cdot]}(\mathbb{S}^1)_+$ (or what is just the same the flag manifold $M(W_\infty^r)$ of the real form W_∞^r of the Pope-Shen-Romans W_∞ -algebra), to consider the squashed product $M(W_\infty^r) \cdot Q^*$ and to justify that W_∞ -algebra acts on $M(W_\infty^r) \cdot Q^*$ by vector fields. To perform this program one needs in a detailed description of the flag manifold $M(W_\infty^r)$.

The flag manifold $M(W_\infty^r)$ may be defined as a symplectic leaf of the Poisson manifold $(W_\infty^r)^*$, the coadjoint module for the Lie algebra W_∞^r (cf.[4]). The tangent space of $M(W_\infty^r)$ at the initial point may be identified with the quotient $W_\infty/W_\infty^{\text{reg}}$, so the flag manifold $M(W_\infty^r)$ is an almost complex manifold.

Proposition 4. *The almost complex structure on the flag manifold $M(W_\infty^r)$ is integrable.*

Proof. First of all, the almost complex structure on $M(W_\infty^r)$ is formally integrable. To prove that it is really integrable one should to construct an almost complex embedding of $M(W_\infty^r)$ into some infinite dimensional complex manifold. The standard manifold for such purposes is an infinite dimensional Grassmannian, f.e. one of subspaces in W_∞ , one of which is W_∞^{reg} .

Being the symplectic leaf of the Poisson manifold $(W_\infty^r)^*$ the flag space $M(W_\infty^r)$ possesses an infinite dimensional family of symplectic structures $\omega_{\mathbf{h},c}$, where c corresponds to the central charge of W_∞^r and $\mathbf{h} = (h_2, h_3, \dots, h_n, \dots)$ corresponds to a character of subalgebra W_∞^{reg} . Coupled with the complex structure 2-forms $\omega_{\mathbf{h},c}$ define an infinite dimensional family of (pseudo)Kähler metrics on the flag manifold $M(W_\infty^r)$.

Each (pseudo)Kähler metric defines a prequantization bundle $E_{\mathbf{h},c}(M(W_\infty^r))$ over $M(W_\infty^r)$, which is a Hermitean line bundle of with the action of the Lie algebra W_∞ by covariant derivatives with curvature form $2\pi\omega_{\mathbf{h},c}$.

In the Fock spaces $F(M(W_\infty^r), E_{\mathbf{h},c})$ there are realized the Verma modules over Pope-Shen-Romans W_∞ algebra (Fock space of a pair (M, E) is a space dual to the space of sections of the bundle E^* over M [13: D.Juriev, Algebra Anal.]). It should be mentioned that Verma modules over $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)$ were investigated recently by V.Kac and A.Radul [9].

2.2. W -symmetries of a second quantized free string (flat background): W -ghosts, W -differential Banks-Peskin forms, Siegel W -string fields and W -BRST-operator in them; W -string Gauss-Manin connection and W -string Kostant-Blattner-Sternberg pairings; geometrical non-cancelation of Bowick-Rajeev W -anomaly — absence of Bowick-Rajeev W -vacua and gauge-invariant Siegel W -string fields; operator cancelation of Bowick-Rajeev W -anomaly — W_N -algebras.

It should be mentioned that the flag manifold $M(W_\infty^r)$ for the W -algebra W_∞^r admits an embedding into the infinite dimensional analogue of the classical symmetric domain of type I [39]. Such embedding is defined by the mapping $W_\infty^r \mapsto \mathfrak{gl}(\infty)$, where $\mathfrak{gl}(\infty)$ is the Lie algebra of linear operators in the space $\mathcal{F}(\mathbb{S}^1)$ of functions on the circle $\mathbb{S}^1$. Because the infinite dimensional classical symmetric domain of type I admits a representation as a space of complex structures on $\mathcal{F}(\mathbb{S}^1) = \text{Map}(\mathbb{S}^1, \mathbb{R})$ as well as on $\text{Map}(\mathbb{S}^1, \mathbb{R}^n)$, then the obtain such representation for the flag manifold $M(W_\infty^r)$, too (such representation is analogous to one of M.Bowick and S.Rajeev for the flag manifold $M(\text{Vir})$ for the Virasoro-Bott group [40]). Thus we have constructed the squash product $M(W_\infty^r) \cdot Q^*$.

Proposition 5. *The mapping $W_\infty^{\text{reg}} \mapsto \mathbb{C}\text{Vect}(M(\text{Vir}) \cdot Q^*)$ is extended to the mapping $W_\infty \mapsto \mathbb{C}\text{Vect}(M(W_\infty^r) \cdot Q^*)$.*

This proposition is a specialization of a standard fact of the theory of inductions to our infinite dimensional case.

Now we should to mention that Nambu-Goto action (the Kähler potential on Q^*) defines the bundle $E_{\text{NG}}(Q^*)$, which may be lifted to the bundle $E_{\text{NG}}(M(W_\infty^r) \cdot Q^*)$; this procedure defines the first cohomology class $H^1(W_\infty; \mathcal{O}(M(W_\infty^r) \cdot Q^*))$ of the Lie algebra W_∞ with coefficients in "classical string fields" $\mathcal{O}(M(W_\infty^r) \cdot Q^*)$ (cf. [1], see also [30]).

The Hermitean line bundle $E_{\text{h},c}(M(W_\infty^r))$ over the flag manifold $M(W_\infty^r)$ may be lifted to the bundle $E_{\text{h},c}(M(W_\infty^r) \cdot Q^*)$ over the space $M(W_\infty^r) \cdot Q^*$, in which the action of the Pope-Shen-Romans algebra W_∞ with non-trivial central charge is defined. One may also include the first cohomology class of W_∞ into such action. That means that one should consider the bundle $\tilde{E}_{\text{h},c}(M(W_\infty^r) \cdot Q^*)$, the tensor product of $E_{\text{h},c}(M(W_\infty^r) \cdot Q^*)$ and $E_{\text{NG}}(M(W_\infty^r) \cdot Q^*)$, instead of $E_{\text{h},c}(M(W_\infty^r) \cdot Q^*)$.

The Fock space $F(M(W_\infty^r) \cdot Q^*), \tilde{E}_{\text{h},c}(M(W_\infty^r) \cdot Q^*)$ over the pair $(M(W_\infty^r) \cdot Q^*), \tilde{E}_{\text{h},c}(M(W_\infty^r) \cdot Q^*)$ is just "the configuration space for a second quantized free string without ghosts after the accounting of W -symmetries" (cf.[1], see also [29]).

Our next purpose is to introduce W -ghosts, to consider W -differential Banks-Peskin forms, Siegel W -string fields, W -BRST-operator in them and, thus, to construct "the configuration space for a second quantized free string with ghosts after the accounting of W -symmetries" in a way analogous to one of [1] (see also [29]).

Unfortunately, this program can not be performed even on the first step. The deal is that one can not construct correctly the semi-infinite forms for the Pope-Shen-Romans W_∞ -algebra (or for $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$). So we should to reduce our geometric picture to a finite order n . Namely, in view of the (real) Radul mapping $\mathcal{F}((W_\infty^r)^*) \mapsto \text{GD}(\mathfrak{sl}(n, \mathbb{R}))$ one may consider the flag manifold $M(\text{GD}(\mathfrak{sl}(n, \mathbb{R})))$ for Gelfand-Dickey algebra instead of the flag manifold $M(W_\infty)$ for W_∞ -algebra. The flag manifold $M(\text{GD}(\mathfrak{sl}(n, \mathbb{R})))$ may be defined as a symplectic leaf of Gelfand-Dickey algebra $\text{GD}(\mathfrak{sl}(n, \mathbb{R}))$. The flag manifold $M(\text{GD}(\mathfrak{sl}(n, \mathbb{R})))$ is a complex man-

ifold as well as $M(W_\infty^r)$ and possesses (at least, one [4]) n -parametric family of symplectic structures $\omega_{\mathbf{h},c}$, where $\mathbf{h} = (h_2, \dots, h_n)$. Coupled with the complex structure these 2-forms define n -parametric family of (pseudo)Kähler metrics on the flag manifold $M(\text{GD}(\text{sl}(n, \mathbb{R})))$; each (pseudo)Kähler metric defines a prequantization bundle $E_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ over $M(\text{GD}(\text{sl}(n, \mathbb{R})))$.

Proposition 6. *The mapping $W_\infty \mapsto \mathbb{C}\text{Vect}(M(W_\infty^r) \cdot Q^*)$ may be reduced to the mapping $W_\infty \mapsto \mathbb{C}\text{Vect}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*)$.*

Proof. It is just the consequence of the existence of Radul mapping.

One may perform for $M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*$ all geometric procedures described above for $M(W_\infty^r) \cdot Q^*$. Namely, the bundle $E_{\text{NG}}(Q^*)$ may be lifted to the bundle $E_{\text{NG}}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*)$; the Hermitean line bundle $E_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ may be lifted to the bundle $E_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*)$; finally, one may consider the bundle $\tilde{E}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*)$ as the tensor product of two previous ones. It is natural consider the Fock space $F(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*, \tilde{E}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*))$ as "the reduced configuration space for a second quantized free string without ghosts after the accounting of W -symmetries".

But before a consideration of the corresponding ghosts it is reasonable to consider the structure of action of the Lie algebra $W_\infty^{\mathbb{C}}$ on $M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*$ more systematically.

It seems that it is very convenient to restrict ourselves (as above) to a consideration of differential operators of order less or equal to n . It means that we shall deal with the quotient $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+ / \widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}$ of the central extension of the Lie algebra $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$ of differential operators without free terms by the central extension of its subalgebra $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1}$ of differential operators without free terms, which are not contained terms with D^k ($1 \leq k \leq n$). The corresponding strict sequence

$$0 \longrightarrow \widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1} \longrightarrow \widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+ \longrightarrow \widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+ / \widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_{\geq n+1} \longrightarrow 0$$

may be resolved. The resolving mapping identifies the quotient with the subspace $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_{+; \leq n}$ of differential operators of order less or equal to n without free terms in the Lie algebra $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$. The object, which elements of $\text{DOP}^{\mathbb{C}}(\mathbb{S}^1)_{+; \leq n}$ form is a Lie quasi(pseudo)algebra, which will be called *enlarged Gervais-Matsuo quasi(pseudo)algebra* and will be denoted $\mathcal{GM}_{n;(\text{enl})}^{\mathbb{C}}$. Gervais-Matsuo quasi(pseudo)algebra may be comprehended as "a less more than one-half of" the enlarged Gervais-Matsuo quasi(pseudo)algebra. It should be mentioned that the enlarged Gervais-Matsuo quasi(pseudo)algebra is embedded in $\widehat{\text{DOP}}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$ rather than in $\text{DOP}_{[\cdot, \cdot]}^{\mathbb{C}}(\mathbb{S}^1)_+$ so we have obtained a central extension of the enlarged Gervais-Matsuo quasi(pseudo)algebra, which will be denoted by $\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}$. If the enlarged Gervais-Matsuo quasi(pseudo)algebra $\mathcal{GM}_{n;(\text{enl})}^{\mathbb{C}}$ acts on $M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*$, then its central extension $\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}$ acts in the line bundles over it.

So now we are able to introduce W -ghosts (related to the central extended enlarged Gervais-Matsuo quasi(pseudo)algebra), to consider the corresponding W -differential Banks-Peskin forms and Siegel W -fields, W -BRST-operator in them

and, thus, to construct "the reduced configuration space for a second quantized free string with ghosts after the accounting of W -symmetries". Namely, W -ghosts may be identified with elements of the Lie quasi(pseudo)algebra $\mathcal{GM}_{n;(\text{enl})}^{\mathbb{C}}$ acting on the manifold $M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*$ by vector fields, W -antighosts are dual 1-forms on $M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*$. W -differential Banks-Peskin forms are just differential forms on the manifold $M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*$ valued in the line bundle $\tilde{E}_{\mathbf{h},c}$ generated by W -antighosts; *semi-infinite W -differential Banks-Peskin forms* are defined with respect to the $\mathbb{Z}$ -grading on the Lie quasi(pseudo)algebra $\mathcal{GM}_{n;(\text{enl})}^{\mathbb{C}}$.

The space of W -differential Banks-Peskin forms should be denoted by $\Omega_{\text{BP}}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*; \tilde{E}_{\mathbf{h},c})$ and the space of semi-infinite W -differential Banks-Peskin forms should be denoted by $\Omega_{\text{BP}}^{\text{SI}}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*; \tilde{E}_{\mathbf{h},c})$.

Remark 1. The Lie quasi(pseudo)algebra $\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}$ acts on the space $\Omega_{\text{BP}}^{\text{SI}}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*; \tilde{E}_{\mathbf{h},c})$ with the central charge $c - 2(2n^3 - n - 1)$.

Definition 2.

1. *Siegel W -string fields* are elements of the space, dual to the space $\Omega_{\text{BP}}^{\text{SI}}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*; \tilde{E}_{\mathbf{h},c})$ of semi-infinite W -differential Banks-Peskin forms.
2. *W -BRST-operator* is the operator Q_{BRST} in the space of Siegel W -string fields dual to the exterior covariant derivative D in the space of semi-infinite W -differential Banks-Peskin forms.

Remark 2. $Q_{\text{BRST}}^2 = 0$, iff $c = 2(2n^3 - n - 1)$.

Let us now study the aspects related to gauge-invariance of W -string fields in sense of [1] (see also [31]).

Namely, the space $\Omega_{\text{BP}}^{\text{SI}}(M(\text{GD}(\text{sl}(n, \mathbb{R}))) \cdot Q^*; \tilde{E}_{\mathbf{h},c})$ of semi-infinite W -differential Banks-Peskin forms may be considered as a space of holomorphic sections of a certain bundle over $M(\text{GD}(\text{sl}(n, \mathbb{R})))$, which will be called *Fock-plus-ghost bundle* and denoted by $\text{FG}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$. Fibers of the vector bundle $\text{FG}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ over points x of the flag manifold $M(\text{GD}(\text{sl}(n, \mathbb{R})))$ for Gelfand-Dickey algebra $\text{GD}(\text{sl}(n, \mathbb{R}))$ will be denoted by $(\text{FG}_{\mathbf{h},c})_x(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$.

There exists a set $\{P_x\}$ of natural gauge-fixing projectors

$$P_x : \mathcal{O}(\text{FG}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R})))) \mapsto (\text{FG}_{\mathbf{h},c})_x(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$$

(here the flag manifold $M(\text{GD}(\text{sl}(n, \mathbb{R})))$ is interpreted as a space of internal gauge degrees of freedom, cf. [31]) and a set $\{I_x\}$ of embedding operators

$$I_x : (\text{FG}_{\mathbf{h},c})_x(M(\text{GD}(\text{sl}(n, \mathbb{R})))) \mapsto \mathcal{O}(\text{FG}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))),$$

which obey two properties;

- (1) $P_x I_x = \text{id}$;
- (2) $\left(\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}\right)_+(x) I_x = 0$, where $\left(\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}\right)_+(x)$ is the natural resolution of the strict sequence

$$\begin{aligned} 0 \longrightarrow \left(\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}\right)_0(x) \longrightarrow \widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}} \longrightarrow \\ \widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}} / \left(\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}\right)_0(x) \longrightarrow 0, \end{aligned}$$

where $\left(\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}\right)_0(x) = \{v \in \widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}} : v(x) = 0\}$.

One may define *W-string Gauss-Manin connection* in $\text{FG}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ as

$$\begin{aligned} \nabla_v \varphi(x) &= \lim_{t \rightarrow \infty} t^{-1} (P_x I_{x+tv(x)} P_{x+tv(x)} \varphi - P_x \varphi), \\ v &\in \text{Vect}(M(\text{GD}(\text{sl}(n, \mathbb{R})))), x \in M(\text{GD}(\text{sl}(n, \mathbb{R}))). \end{aligned}$$

This connection may be also define by the use of *W-string Kostant-Blattner-Sternberg pairings*. Namely, the Fock space $F(M(\text{GD}(\text{sl}(n, \mathbb{R}))), \text{FG}_{\mathbf{h},c})$ possesses a (pseudo)hermitean metric (cf.[13:D.Juriev, Algebra Anal.,31]). If such metric is non-degenerate then it induces a metric $(\cdot, \cdot)$ in the space $\mathcal{O}(\text{FG}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$. *W-string Kostant-Blattner-Sternberg pairings* are the mappings $B_{x,y}(\cdot, \cdot)$ from the tensor product of $(\text{FG}_{\mathbf{h},c})_x(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ and $(\text{FG}_{\mathbf{h},c})_y(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ to $\mathbb{C}$, such that $B_{x,y}(\varphi, \psi) = (I_x \varphi, I_y \psi)$.

The *W-string Gauss-Manin connection* ∇ may be expressed via *W-string Kostant-Blattner-Sternberg pairings* in the following way:

$$\begin{aligned} \nabla_v \Phi(x) &= 0 \text{ iff } B_{x+tv(x),x}(\Phi(x+tv(x)), \psi) = B_{x,x}(\Phi(x), \psi) + o(t) \\ &\text{for all } \psi \in (\text{FG}_{\mathbf{h},c})_y(M(\text{GD}(\text{sl}(n, \mathbb{R})))), \end{aligned}$$

where $\Phi(x)$ is a short notation for $P_x(\Phi)$.

Definition 3. (cf. [31]).

1. A covariantly constant section of Fock-plus-ghost bundle $\text{FG}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ over the flag manifold $M(\text{GD}(\text{sl}(n, \mathbb{R})))$ for Gel'fand-Dickey algebra $\text{GD}(\text{sl}(n, \mathbb{R}))$ is called *Bowick-Rajeev W-vacuum*.
2. The space dual to the space of Bowick-Rajeev vacua is called *the space of gauge-invariant Siegel W-string fields*.

Unfortunately, Bowick-Rajeev W -vacua (or equivalently gauge-invariant Siegel W -string fields) do not exist. *The phenomenon of geometric non-cancellation of Bowick-Rajeev W -anomaly may be considered as an explanation of the fact that Gelfand-Dickey brackets can not be quantized only by addition of central term.* The global structural change of commutation relations is necessary. Let us now describe the process of *operator cancelation* of Bowick-Rajeev anomaly, which transforms Gelfand-Dickey algebras or enlarged Gervais-Matsuo quasi(pseudo)algebras into W_N -algebras (it should be mentioned that problem of quantization of Lie quasi(pseudo)algebras was discussed in another context in the book of M.V.Karasev and V.P.Maslov [33]). Of course, such operator cancelation does not supply us with BRST-operator for W_N -algebras.

Let us consider an arbitrary point x of the flag manifold $M(\text{GD}(\text{sl}(n, \mathbb{R})))$ for Gelfand-Dickey algebra $\text{GD}(\text{sl}(n, \mathbb{R}))$. Let us embed the fiber $(\text{FG}_{\mathbf{h},c})_x(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ of Fock-plus-ghost bundle $\text{FG}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ over x into the space $\mathcal{O}(\text{FG}_{\mathbf{h},c}(M(\text{GD}(\text{sl}(n, \mathbb{R}))))$ by use of I_x . Now define the action of the real form $\widehat{\mathcal{GM}}_{n;(\text{enl})}$ of enlarged Gervais-Matsuo quasi(pseudo)algebra $\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}$ in V in the following way:

$$v(\phi) = P_x \nabla_v I_x \phi, \quad \phi \in (\text{FG}_{\mathbf{h},c})_x(M(\text{GD}(\text{sl}(n, \mathbb{R})))), v \in \widehat{\mathcal{GM}}_{n;(\text{enl})}.$$

It is not the action indeed because the commutation relations in the real form of Gervais-Matsuo quasi(pseudo)algebra are broken, moreover, the object, which we have received in no more a Lie quasi(pseudo)algebra, but an ordinary algebra of operators.

It is just remarkable that the obtained algebras are just (after a complexification) W_N -algebras of papers [3].

CONCLUSIONS

Thus, our plan is performed, the results may be summarized.

It was appeared that the main objects of the infinite dimensional W -geometry of a second quantized free string are not infinite dimensional groups, Lie algebras and their homogeneous spaces as it was in [1] but infinite dimensional Lie quasi(pseudo)algebras (various modifications of Gervais-Matsuo quasi(pseudo)-algebra $\mathcal{GM}_n^{\mathbb{C}}$ of classical (infinitesimal) W -transformations), nonlinear Poisson brackets and related geometrical structures.

As a consequence there exists a geometrical non-cancellation of Bowick-Rajeev anomaly (the absence of gauge-invariant Siegel W -string fields). Operator cancelation provides us with transformation of classical Lie quasi(pseudo)algebras $\widehat{\mathcal{GM}}_{n;(\text{enl})}^{\mathbb{C}}$ (central extended enlarged Gervais-Matsuo algebras) into quantum W_N -algebras.

It should be mentioned that because realistic W -string field theory is essentially the theory of self-interacting string field (see f.e. [41]), free W -string field theory may be considered only as a startpoint of the perturbation approach to it. Respectively, the infinite dimensional W -geometry of a second quantized free string may be also regarded as a zero-approximation of the noncommutative geometry of self-interacting W -string field.

REFERENCES

- [1] D.Juriev, Infinite dimensional geometry and quantum field theory of strings. I. Infinite dimensional geometry of a second quantized free string, *Alg. Groups Geom.* 11 (1994).
- [2] D.Juriev, Infinite dimensional geometry and quantum field theory of strings. II. Infinite dimensional noncommutative geometry of self-interacting string field, [in preparation].
- [3] A.B.Zamolodchikov, Infinite additional symmetries in 2-dimensional conformal field theory, *Theor. Math. Phys.* 65 (1985) 1205-1213;
A.B.Zamolodchikov and V.A.Fateev, Conformal quantum field theory in two dimensions having $\mathbb{Z}_3$ -symmetry, *Nucl. Phys. B* 280 (1987) 644-660;
S.L.Lukyanov and V.A.Fateev, Conformally invariant models in 2-dimensional quantum field theory with $\mathbb{Z}_N$ symmetry, *Zh. Exp. Theor. Fiz.* 94(3) (1988) 23-37;
S.L.Lukyanov, Quantization of the Gel'fand-Dickey bracket, *Funkt. Anal. Appl.* 22 (1988) 255-263;
J.-L.Gervais, Systematic approach to conformal theories, *Nucl. Phys. B (Proc. Suppl.)* 5B (1988) 119-136;
J.-L.Gervais and A.Bilal, Conformal theories with non-linearly-extended Virasoro symmetries and Lie algebra classification. In: *Infinite dimensional Lie algebras and Lie groups* Ed. V.Kac (World Scientific, Singapore, 1989) 483-526; Systematic approach to conformal field systems with extended Virasoro symmetries, *Phys. Lett. B* 206 (1988) 412-420; Systematic construction of conformal theories with higher-spin Virasoro symmetries, *Nucl. Phys. B* 318 (1989) 579-630.
- [4] V.Yu.Ovsienko and O.D.Ovsienko, Lie derivatives of order n on the line. Tensor meaning of the Gel'fand-Dickey bracket, *Adv. Soviet Math.* 2 (1991) 221-231;
V.Yu.Ovsienko and B.A.Khesin, Symplectic leaves of the Gel'fand-Dickey brackets and homotopy classes of nondegenerate curves, *Funkt. Anal. Appl.* 24 (1990) 33-40.
- [5] O.S.Kravchenko and B.A.Khesin, A central extension of the algebra of pseudodifferential symbols, *Funkt. Anal. Appl.* 25 (1991) 152-154.
- [6] A.O.Radul, Lie algebras of differential operators, their central extensions and W -algebras, *Funkt. Anal. Appl.* 25 (1991) 33-49; Non-trivial central extensions of Lie algebras of differential operators in two and higher dimensions, *Phys. Lett. B* 265 (1991) 86-91.
- [7] I.Vaysburd and A.Radul, Differential operators and W -algebra, *Phys. Lett. B* 274 (1992) 317-322.
- [8] I.Bakas, B.Khesin and E.Kiritsis, The logarithm of the derivative operator and higher spin algebras of W_∞ type, *Commun. Math. Phys.* 151 (1993) 233-243.
- [9] V.Kac and A.Radul, Quasilinear highest weight modules over the Lie algebra of differential operators on the circle, *Commun. Math. Phys.* 157 (1993) 429-458.
- [10] B.Khesin and I.Zakharevich, Poisson-Lie group of pseudodifferential operators and fractional KP-KdV hierarchies, *C. R. Acad. Sci. Paris I*, 316 (1993) 621-626.
- [11] J.-L.Gervais and Y.Matsuo, W -geometries, *Phys. Lett. B* 274 (1992) 309-316; Classical A_n - W -geometry, *Commun. Math. Phys.* 152 (1993) 317-368;
J.-L.Gervais, W -geometry from chiral embeddings, *J. Geom. Phys.* 11 (1993) 293-304; Introduction to differential W -geometry, preprint LPTENS-93/38.
- [12] P.Griffiths and S.Harris, *Principles of Algebraic Geometry* (Wiley Interscience, New York, 1978).
- [13] A.A.Kirillov, A Kähler structure on K -orbits of the group of diffeomorphisms of a circle, *Funkt. Anal. Appl.* 21 (1987) 122-125;
A.A.Kirillov and D.V.Juriev, The Kähler geometry on the infinite dimensional homogeneous manifold $M = \text{Diff}_+(\mathbb{S}^1)/\text{Rot}(\mathbb{S}^1)$, *Funkt. Anal. Appl.* 20 (1986) 322-324; The Kähler geometry on the infinite dimensional homogeneous space $M = \text{Diff}_+(\mathbb{S}^1)/\text{Rot}(\mathbb{S}^1)$ *Funkt. Anal. Appl.* 21 (1987) 284-293; Representations of the Virasoro algebra by the orbit method, *J. Geom. Phys.* 5 (1988) 351-364 [reprinted in *Geometry and Physics. Essays in honour of I.M.Gelfand*. Eds. S.G.Gindikin and S.G.Singer (Pitagora Editrice, Bologna and Elsevier Sci. Publ., Amsterdam, 1991)];
D.Juriev, The vocabulary of geometry and harmonic analysis on the infinite-dimensional manifold $\text{Diff}_+(\mathbb{S}^1)/\mathbb{S}^1$, *Adv. Soviet Math.* 2 (1991) 233-247; A model of Verma modules over the Virasoro algebra, *Algebra Anal.* 2(2) (1990) 209-226; An infinite dimensional geometry of the universal deformation of a complex disc. *Russian J. Math. Phys.* 2(1)

- (1994).
- [14] D.Juriev, Quantum conformal field theory as infinite dimensional noncommutative geometry, Russian Math.Surveys 46(4) (1991) 135-163.
 - [15] A.Yu.Morozov, Strings – what are they? Uspekhi Fiz. Nauk 162(8) (1992) 84-175; 162(11) (1992) 206.
 - [16] D.Juriev, Quantum projective field theory: quantum field analogs of Euler formulas, Teor. Matem. Fiz. 92(1) (1992) 172-176; Quantum projective field theory: quantum field analogs of Euler-Arnold equations in projective G -hypermultiplets, Teor. Matem. Fiz. (1994).
 - [17] V.I.Arnold, *Mathematical methods of classical mechanics* (Springer, 1976).
 - [18] E.Floratos and J.Iliopoulos, A note on the classical symmetries of the closed bosonic membranes, Phys. Lett. B 201 (1988) 237-240;
I.Antoniadis, P.Ditsas, E.Floratos and J.Iliopoulos, New realizations of the Virasoro algebra as membrane symmetries, Nucl. Phys. B 300 (1988) 549-558.
 - [19] I.M.Gelfand and D.B.Fairlie, The algebra of Weyl symmetrized polynomials and its quantum extension, Commun. Math. Phys. 136 (1991) 487-499.
 - [20] D.B.Fairlie, P.Fletcher and C.K.Zachos, Trigonometric structure constants for new infinite-dimensional algebras, Phys. Lett. B 218 (1989) 203-206;
D.B.Fairlie and C.K.Zachos, Infinite dimensional algebras, sine brackets and $SU(\infty)$, Phys.Lett. B 224 (1989) 101-107;
D.B.Fairlie, C.K.Zachos and P.Fletcher, Infinite dimensional algebras and a trigonometric basis for the classical Lie algebras, J. Math. Phys. 31 (1990) 1088-1094.
 - [21] V.Kac and D.H.Peterson, Spin and wedge representations of infinite-dimensional Lie algebras and groups, Proc. Nat'l Acad. Sci. USA 78 (1981) 3308-3312.
 - [22] D.B.Fairlie and J.Nuyts, Deformations and renormalisations of W_∞ , Commun. Math. Phys. 134 (1990) 413-419.
 - [23] I.Bakas, The large- N limit of extended conformal symmetries, Phys. Lett. B 228 (1989) 57-63.
 - [24] C.N.Pope, X.Shen and L.J.Romans, The complete structure of W_∞ , Phys. Lett. B 236 (1990) 173-178; A new higher-spin algebra and the lone-star product, Phys. Lett. B 242 (1990) 401-406; W_∞ and the Racah-Wigner algebra, Nucl. Phys. B 339 (1990) 191-221.
 - [25] I.Bakas, The structure of the W_∞ -algebra, Commun. Math. Phys. 134 (1990) 487-508.
 - [26] I.N.Bernstein, I.M.Gelfand and S.I.Gelfand, Models of representations of Lie groups. Trudy Semin. I.G.Petrovskogo 2 (1976) 3-21.
 - [27] B.L.Feigin, The Lie algebras $gl(\lambda)$ and the cohomology of the Lie algebra of differential operators, Russian Math. Surveys 35(2) (1988) 169-170.
 - [28] D.V.Juriev, Complex projective geometry and quantum projective field theory, Teor. Matem. Fiz. (submitted).
 - [29] D.Juriev, Infinite dimensional geometry and closed string-field theory, Lett. Math. Phys. 22 (1991) 1-6.
 - [30] D.Juriev, Cohomology of the Virasoro algebra with coefficients in string fields. Lett. Math. Phys. 19 (1990) 355-356.
 - [31] D.Juriev, Gauge invariance of Banks-Peskin differential forms (flat background), Lett. Math. Phys. 19 (1990) 59-64.
 - [32] I.Batalin, Quasigroup construction and first class constraints, J. Math. Phys. 22 (1981) 1837-1850.
 - [33] M.V.Karasev and V.P.Maslov, *Nonlinear Poisson brackets. Geometry and quantization* (Nauka, Moscow, 1991).
 - [34] L.V.Sabinin, On the equivalence of the category of loops and the category of homogeneous spaces, Soviet Math. 13 (1972) 533-536.
 - [35] P.O.Mikheev and L.V.Sabinin, On the infinitesimal theory of local analytic loops. Soviet Math. 36 (1988) 545-548.
 - [36] J.-L.Gervais and M.V.Saveliev, W -geometry of the Toda systems associated with non-exceptional simple Lie algebras, preprint LPTENS-93/47.
 - [37] P.O.Mikheev and L.V.Sabinin, Smooth quasigroups and geometry. VINITI Problems in Geometry 20 (1988) 75-110 [and several unpublished texts on "nonlinear geometric algebra"].
 - [38] Yu.I.Manin, *Cubic forms* (Nauka, Moscow, 1969).

- [39] E.Cartan, Sur les domaines bornés homogènes de l'espace de n variables complexes, Abh. Math. Seminar Hamburg 11 (1935) 116-162 [reprinted in *Oeuvres Completes* (Gauthier-Villars, Paris, 1953)].
- [40] M.J.Bowick and S.G.Rajeev, String theory as the Kähler geometry of loop space, Phys. Lett. Rev. 58 (1987) 535-538; The holomorphic geometry of closed bosonic string theory and $\text{Diff}(\mathbb{S}^1)/\mathbb{S}^1$, Nucl. Phys. B 293 (1987) 348-384.
- [41] A.Bilal and J.-L.Gervais, Non-linearly extended Virasoro algebras: new prospects for building string theories, Nucl. Phys. B 326 (1989) 222-236.